

# Solitary electromechanical pulses in Lobster neurons.

Alfredo Gonzalez-Perez<sup>1</sup>, Lars D. Mosgaard<sup>1</sup>, Rima Budvytyte<sup>1</sup>, Edgar Villagran-Vargas<sup>3</sup>, Andrew D. Jackson<sup>2</sup> and Thomas Heimburg<sup>1,\*</sup>

<sup>1</sup>Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, 2100 Copenhagen Ø, Denmark

<sup>2</sup>Niels Bohr International Academy, University of Copenhagen, Blegdamsvej 17, 2100 Copenhagen, Denmark

<sup>3</sup>Facultad de Ciencias, Universidad Autónoma del Estado de México. Instituto Literario 100, Colonia Centro, 50000, Toluca, México

**ABSTRACT** Investigations of nerve activity have focused predominantly on electrical phenomena. Nerves, however, are thermodynamic systems, and changes in temperature and in the dimensions of the nerve can also be observed during the action potential. Measurements of heat changes during the action potential suggest that the nerve pulse shares many characteristics with an adiabatic pulse. In the past, experiments by Iwasa & Tasaki demonstrated small changes in nerve thickness and length during the action potential. Such findings have led to the suggestion that the action potential may be related to electromechanical solitons traveling without dissipation. Here, we present ultrasensitive AFM recordings of mechanical changes on the order of 0.2 – 1.2 nm in the giant axons of the lobster. When stimulated at opposite ends of the same axon, colliding action potentials pass through one another and do not annihilate. These observations are consistent with a mechanical interpretation of the nervous impulse.

**Keywords:** Nerves, thermodynamics, hydrodynamics, atomic force microscopy, electrophysiology, solitons

\*corresponding author, theimbu@nbi.dk.

## Introduction

It is not generally appreciated that the size of the nervous impulse is remarkably large. A myelinated motor neuron has a pulse velocity on the order of 100 m/s. Given a typical pulse duration of 1 ms, the resulting size of the nerve pulse is 10 cm. Propagation velocities in slow, non-myelinated fibers are reduced to 1-10 m/s, indicating a pulse size of 1-10 mm. Thus, nerve pulses are macroscopic phenomena spanning a significant fraction of the total axon length. In some cases they can even be larger than small neurons such as interneurons that are only a few 100  $\mu\text{m}$  long (1). In the past, the activity of nerves has conventionally been considered to be a purely electrical phenomenon produced by the flux of ions and the charging of the membrane capacitor (2). The shear size of a nerve pulse suggests that the macroscopic thermodynamic properties of the nerve membrane or of the entire nerve ought to be taken into account. It is to be expected that the state of the nerve cell depends not only on electrochemical potentials and the conjugated flux of ions but also on all other thermodynamic forces including variations in lateral pressure (resulting in changes of membrane area and thickness) and temperature (resulting in heat flux). It is therefore not surprising that, during the action potential, one finds changes not only in voltage but also in thickness (3–6), length (5, 7) as well changes in membrane temperature (8–11). The change of thickness of a single squid axon was found to be on the order of 1 nm, and temperature changes range between 1–100  $\mu\text{K}$  depending on the specimen. While both mechanical and thermal signals are very small, they are found to be in phase with voltage changes. The heat signal can be blocked by neurotoxins such as tetrodotoxin (11). This strongly suggests that these thermodynamic phenomena are correlated with the voltage pulse and do not represent independent secondary phenomena.

From a thermodynamic viewpoint, one can consider two extreme cases of possible dynamic changes associated with the action potential: 1. Purely dissipative processes during which entropy increases, such as the flow of ions along concentration gradients. Such processes form the basis of the Hodgkin-Huxley model for the action potential. 2. Adia-

batic processes that do not dissipate heat and thus conserve entropy. These phenomena are rather governed by the laws of analytical mechanics. They play an important role for dynamic properties such as sound propagation. In this context, the heat changes observed in nerves are of fundamental interest. It was found that heat is released during an initial phase of the action potential and that this heat is reabsorbed in the final phase of the action potential. During the nerve pulse no heat (or only very little) is dissipated, and the entropy of the membrane is basically conserved (9, 11). Thus, thermal measurements suggest that the action potential is an adiabatic phenomenon reminiscent of a sound wave. As early as 1912, the striking absence of heat production led Hill to conclude: ‘*This suggests very strongly ... that the propagated nervous impulse is not a wave of irreversible chemical breakdown, but a reversible change of a purely physical nature*’ (12). In contrast, the contemporary understanding of the nerve pulse is based on the flow of ions along gradients through ion channel proteins (2) and therefore assumes that it is of a dissipative nature. Hodgkin himself compared it to the ‘*burning of a fuse of gunpowder*’ (13). There is thus a disagreement between electrophysiological models and some thermodynamics findings. These problems are not easily resolved and merit careful attention.




Figure 1: Schematic representation of a mechanical soliton traveling in a nerve fiber. Small local changes in thickness are caused by pressure-induced order transitions in the membrane. Red regions correspond to ordered lipids in the otherwise liquid (green) lipid membrane. In a living nerve, the spatial extension of the pulse is much larger than shown here.




Figure 2: *Outline of the mechanical experiment. A. Image of the two lobster connectives including the brain (top) and the subesophageal ganglion (bottom). B. Detail of a lobster connective with the lateral giant axon exposed. C. Schematic representation of the cross-section of a lobster connective containing a medial giant axon, a lateral giant axon and several small fibers. The sheath surrounding the connective is cut in the longitudinal direction. D. After opening the sheaths, the two giant axons are exposed to the outside solution. Mechanical signals are recorded with a tipless AFM cantilever. The open connective is placed on a cell containing several pairs of electrodes to stimulate the nerve and to record the signal. The mechanical response is measured on a planar support in the middle of the cell.*

The observed reversible change in temperature as well as mechanical changes seen in optical and mechanical experiments led to the proposal that the action potential is a consequence of an electromechanical pulse or soliton (14). A condition for the existence of such a soliton is the existence of an order transition in the membrane from solid to liquid slightly below physiological temperature. Such transitions have been found in various biological membranes (15). It is thought that the soliton consists of a region of ordered lipid membrane traveling in the otherwise liquid membrane with a speed somewhat less than the speed of sound in the membrane (14, 16). This is shown schematically in Fig. 1. The difference in membrane thickness between the solid and the liquid is of order 1 nm ( $\approx 17\%$  of the total membrane thickness). The associated reversible change in energy is related to the latent heat of the membrane transition and thus to the reversible heat production found in nerves. Since the membrane changes its thickness, changes in membrane voltage of order 50mV are to be expected as a consequence of changes in its capacitance (17, 18). Thus, the soliton is of an electromechanical or piezoelectric nature. An intrinsic feature of such solitons is that two colliding pulses pass through each other without dissipation (16) rather than annihilating as expected in Hodgkin-Huxley-like pulses due to the refractory period. In fact, the penetration of colliding nerve pulses was seen recently in nerves from earth worm and the ventral cord of lobster (19).

Changes in voltage, thickness and temperature in a soliton are all associated with a single phenomenon, and it is not appropriate to consider some of these changes as side effects of another dominant process. They are rather different aspects of the same phenomenon as seen by different instrumentation. There exists clear evidence that electromechanical pulses can travel on lipid monolayers close to the LE-LC transition with velocities very close to those of non-myelinated

nerves (20, 21). However, there is a striking lack of experiments that actually demonstrate the mechanical nature of the nerve pulse. For this reason, the thermodynamic and mechanical interpretation of the nerve pulse has been widely ignored.

## Mechanical changes

Here, we present the results of experiments on the mechanical changes in single axons from lobster connectives. The ventral cord of the lobster is made of two connective strands containing one lateral and one median giant axons each (LG and MG axons, respectively). Additionally, they contain several small nerve fibers. The action potentials of the giant axons can be clearly distinguished from those of the small fibers. The latter yield much smaller signals and are excited only at higher voltages. The MG axon displays a larger peak amplitude and a larger conduction velocity as compared to the LG axon.

We recorded the mechanical response of the giant axons using atomic force microscopy (AFM). Since the vertical displacement of the axonal membrane is expected to be small, the surface of individual giant axons must be accessed directly by the AFM cantilever. In order to achieve this, the sheath surrounding the connective was cut open. This provides direct access to the single axons (a single exposed LG axon is shown in Fig. 2 B together with the opened connective). The experimental setup is described schematically in Figs. 2C and D, and described in detail in Supplement S1. In brief, a tipless cantilever is placed on a single axon exposed from the opened connective. The open connective is placed on top of a cell with 21 electrodes that allow us to stimulate the nerve and to measure the electrical response. In the center of the cell, the nerve is placed on a support for the mechanical measurement. We stimulated the nerve periodically with a pair of electrodes at one end. We monitor only the vertical displacements




Figure 3: *Vertical displacements as recorded by AFM. Left: Differential voltage change (A), integrated signal (B) and corresponding mechanical displacement in the same neuron (C). Right: Mechanical recordings from six different preparations all yield mechanical changes between 0.2 and 1.2 nm. All recordings have a similar shape with small differences due to the size of the nerve and the precise positioning of the AFM cantilever. Mechanical and electrical signals were not measured at exactly the same location. In this figure, they have been temporally aligned.*

ment of the cantilever without scanning in the  $x$ - $y$  plane. The sampling rate is 40 kHz, and the response time of the AFM setup is about 1 ms. The experimental results are shown in Fig. 3 (left panels). The nerve signal is recorded by two electrodes separated by a distance less than the width of the nerve pulse. Thus, in effect, the first derivative of the true voltage change is measured (Fig. 3A). The integral of this signal is roughly proportional to the true voltage change (Fig. 3B). The bottom trace is the AFM signal (Fig. 3C). In order to achieve a good signal-to-noise ratio we averaged over approximately 100 individual action potentials, excited at intervals of 0.2 seconds. In the experiment shown in Fig. 3C, the vertical displacement was about 2 Å. Fig. 3D) shows mechanical signals from six different lobster specimens. We find displacements between 2 and 12 Å lasting between  $\sim$ 2 and 4 ms. Thus, the mechanical changes in these neurons can be measured in a consistent and reproducible manner. We found these mechanical signals consistently in all connective preparations studied. Electrical recordings demonstrated that more than one single axon was stimulated in the recordings with larger voltage amplitudes. The stimulation artifact visible in the electrical recordings had no influence on the mechanical recording.

## Pulse collision

In a recent publication we showed that action potentials traveling in opposite direction in some nerves (earth worm, lobster ventral cords) can pass through each other upon collision (19). Whether this is a generic feature of nerves is not

clear. In a very early experiment Tasaki showed that colliding pulses in single axons from sciatic nerves of toads annihilate (22). Penetration of pulses speak in favor of a mechanical mechanism for nerve pulse propagation. In contrast, annihilation favors a dissipative ion-channel based view.

Here, we present evidence for pulse penetration in single axons from the connectives that were used in the mechanical experiment. We stimulated an action potential in orthodromic and in antidromic directions. We performed experiments on about 20 lobsters. The voltage was chosen such that only one axon was excited. In about 6 experiments, we severed one axon of the connective in order to eliminate the possibility that pulses in opposite directions traveled in different axons (indicated in Fig. 4, top left). In an additional 3 experiments, we cut one axon at two different locations (indicated in Fig. 4, top right, see supplement S1 for details). In these experiments, there exists only one giant axon between the three pairs of electrodes. Stimulation was made using pairs of electrodes at the ends of the axon. Propagating pulses were recorded with a pair of electrodes at a position closer to the antidromic side (indicated in Fig. 4, top left) or closer to the orthodromic side (indicated in Fig. 4, top right). The intact giant axon (the MG fiber) was stimulated in the orthodromic direction (trace A), in the antidromic direction (trace B) and in both directions (trace C). When increasing the voltage it became evident that only one giant axon could be stimulated. Some residual activity from the small fibers could sometimes be seen. In the experiment shown in Fig. 4, left), the antidromic signal arrives at the recording site prior to the orthodromic signal. When stimulated at both ends, the resulting trace looked similar to the sum of single orthodromic and single antidromic pulses indicating that the pulses penetrated without major perturbation. This is consistent with previous findings in worm axons and other axons in lobster (19). In the experiment shown in Fig. 4, right, the antidromic signal arrives at the recording site after the orthodromic signal. After collision, both pulses can still be seen, indicating that they passed through each other.

## Conclusion

In his textbook from 1964 Alan F. Hodgkin recommended: *"In thinking about the physical basis of the action potential perhaps the most important thing to do at the present moment is to consider whether there are any unexplained observations which have been neglected in an attempt to make the experiments fit into a tidy pattern"* (13). Hodgkin was especially concerned about the temperature changes measured during the action potential (8). While Hodgkin's concerns reflect good scientific praxis, neither the measurement of heat nor mechanical changes associated with the action potential have received the attention that they merit. Here, we have demonstrated that mechanical signals in individual axons from lobster connectives propagate in phase with electrical signals. This is in agreement with very early recordings in squid by Iwasa and Tasaki (3, 4). The vertical amplitudes vary from 0.2-1.2 nm. We have also shown that pulses traveling in opposite directions pass through each other without significant distortion. This is expected for electromechanical pulses or






Figure 4: Pulse collision experiment in the median axon of the lobster connective. The top panels represent a schematic drawing of the connective, the position of the electrodes and of the cuts in the lateral giant axon. Left: The lateral giant axon was cut at one position on the orthodromic side of the recording electrodes. Trace A. The nerve was stimulated in orthodromic direction with the two red electrodes. The signal was recorded with the two blue electrodes. Trace B. The nerve was stimulated in antidromic direction with the two green electrodes and recorded with the blue electrodes. Trace C. The nerve was stimulated from both sides. Both pulses can be measured with the blue electrodes, suggesting that they did not annihilate upon collision. The dashed line is the sum of the orthodromic and the antidromic signals from traces A and B and is given for comparison. Right: A similar experiment in which the lateral giant axon was cut at two positions left and right of the recording electrodes. In this experiment, both orthodromic and antidromic pulses could be seen at the recording electrodes suggesting that they did not annihilate upon collision. The grey-shaded regions display the stimulation artifact. For experimental details see also (19).

solitons. It is, however, unexpected within the framework of a Hodgkin-Huxley mechanism (see also (19)). While these findings do not prove any particular mechanism for the action potentials, they do suggest that it is advisable to study changes in thermodynamic variables in addition to voltage and current.

Recently, there have been various reports, both theoretical and experimental, regarding the possibility of mechanical pulse propagation in artificial systems close to transitions and in nerves (14, 16, 17, 20, 21, 23–25). Heimbürg and Jackson (14) argued that, close to the phase transitions found in biological tissue, electromechanical solitons with properties similar to those of the action potential can travel along the nerve axons. El Hady and Machta (26) recently argued that, in general, the change of charge on the membrane capacitor leads to electrostrictive forces that must alter the membrane dimensions (see also (18)). In (27) it was shown that such an electromechanical picture can provide important insights regarding the mechanism of general anesthesia, which is then seen as the familiar physical chemical effect of anesthetics on melting transitions in biomembranes. It seems likely that the neglect of non-electrical effects has been motivated by the relatively narrow and exclusively electrical focus of the accepted framework provided by Hodgkin and Huxley (2). The study of the mechanical nature of the nerve pulse promises important insights into the underlying thermodynamic nature of the action potential and its control by thermodynamic vari-

ables distinct from voltage.

**Acknowledgments:** This work was supported by the Villum foundation (VKR 022130).

## References

1. Goodman, C. 1974. Anatomy of locust ocellar interneurons - constancy and variability. *J. Comp. Physiol.* 95:185–201.
2. Hodgkin, A. L., and A. F. Huxley. 1952. A quantitative description of membrane current and its application to conduction and excitation in nerve. *J. Physiol. London* 117:500–544.
3. Iwasa, K., and I. Tasaki. 1980. Mechanical changes in squid giant-axons associated with production of action potentials. *Biochem. Biophys. Research Comm.* 95:1328–1331.
4. Iwasa, K., I. Tasaki, and R. C. Gibbons. 1980. Swelling of nerve fibres associated with action potentials. *Science* 210:338–339.
5. Tasaki, I., K. Kusano, and M. Byrne. 1989. Rapid mechanical and thermal changes in the garfish olfactory nerve associated with a propagated impulse. *Biophys. J.* 55:1033–1040.
6. Kim, G. H., P. Kosterin, A. Obaid, and B. M. Salzberg. 2007. A mechanical spike accompanies the action potential in mammalian nerve terminals. *Biophys. J.* 92:3122–3129.
7. Wilke, E., and E. Atzler. 1912. Experimentelle Beiträge zum Problem der Reizleitung im Nerven. *Pflügers Arch.* 146:430–446.

8. Abbott, B. C., A. V. Hill, and J. V. Howarth. 1958. The positive and negative heat production associated with a nerve impulse. *Proc. Roy. Soc. Lond. B* 148:149–187.
9. Howarth, J. V., R. Keynes, and J. M. Ritchie. 1968. The origin of the initial heat associated with a single impulse in mammalian non-myelinated nerve fibres. *J. Physiol.* 194:745–793.
10. Howarth, J. 1975. Heat production in non-myelinated nerves. *Phil. Trans. Royal Soc. Lond.* 270:425–432.
11. Ritchie, J. M., and R. D. Keynes. 1985. The production and absorption of heat associated with electrical activity in nerve and electric organ. *Quart. Rev. Biophys.* 18:451–476.
12. Hill, A. V. 1912. The absence of temperature changes during the transmission of a nervous impulse. *J. Physiol. London* 43:433–440.
13. Hodgkin, A. L., 1964. The conduction of the nervous impulse. Liverpool University Press, Liverpool, UK.
14. Heimburg, T., and A. D. Jackson. 2005. On soliton propagation in biomembranes and nerves. *Proc. Natl. Acad. Sci. USA* 102:9790–9795.
15. Heimburg, T., 2007. Thermal biophysics of membranes. Wiley VCH, Berlin, Germany.
16. Lautrup, B., R. Appali, A. D. Jackson, and T. Heimburg. 2011. The stability of solitons in biomembranes and nerves. *Eur. Phys. J. E* 34:57.
17. Heimburg, T., and A. D. Jackson. 2007. On the action potential as a propagating density pulse and the role of anesthetics. *Biophys. Rev. Lett.* 2:57–78.
18. Heimburg, T. 2012. The capacitance and electromechanical coupling of lipid membranes close to transitions. the effect of electrostriction. *Biophys. J.* 103:918–929.
19. Gonzalez-Perez, A., R. Budvytyte, L. D. Mosgaard, S. Nissen, and T. Heimburg. 2014. Penetration of action potentials during collision in the median and lateral giant axons of invertebrates. *Phys. Rev. X* 4:031047.
20. Griesbauer, J., S. Bössinger, A. Wixforth, and M. F. Schneider. 2012. Propagation of 2d pressure pulses in lipid monolayers and its possible implications for biology. *Phys. Rev. Lett.* 108:198103.
21. Griesbauer, J., S. Bössinger, A. Wixforth, and M. F. Schneider. 2012. Simultaneously propagating voltage and pressure pulses in lipid monolayers of pork brain and synthetic lipids. *Phys. Rev. E* 86:061909.
22. Tasaki, I. 1949. Collision of two nerve impulses in the nerve fiber. *Biochim. Biophys. Acta* 3:494–497.
23. Kaufmann, K., 1989. Lipid membrane. [http://membranes.nbi.dk/Kaufmann/pdf/Kaufmann\\_book5.org.pdf](http://membranes.nbi.dk/Kaufmann/pdf/Kaufmann_book5.org.pdf), Caruaru.
24. Shrivastava, S., and M. F. Schneider. 2014. Evidence for two-dimensional solitary sound waves in a lipid controlled interface and its implications for biological signalling. *J. R. Soc. Interface* 11:20140098.
25. Shrivastava, S., K. H. Kang, and M. F. Schneider. 2015. Solitary shock waves and adiabatic phase transition in lipid interfaces and nerves. *Phys. Rev. E* 91:012715.
26. El Hady, A., and B. B. Machta. 2014. Mechanical surface waves accompany action potential propagation. [arXiv:1407.7600v2 \[q-bio.NC\]](https://arxiv.org/abs/1407.7600v2)
27. Heimburg, T., and A. D. Jackson. 2007. The thermodynamics of general anesthesia. *Biophys. J.* 92:3159–3165.
28. Evans, P. D., E. A. Kravitz, B. R. Talamo, and B. G. Wallace. 1976. The association of octopamine with specific neurones along lobster nerve trunks. *The Journal of physiology* 262:51–70.
29. Wright, E. B., and J. P. Reuben. 1958. A comparative study of some excitability properties of the giant axons of the ventral nerve cord of the lobster, including the recovery of excitability following an impulse. *J. Cell Comp. Physiol.* 51:13–28.

## Supplement 1: Materials and methods

**Materials.** In our experiments we used lobster, *Homarus americanus*, that was obtained from a local supplier that imported the animals from Canada. We used a lobster saline solution adapted from Evans et al. (28) 462 mM NaCl, 10 mM KCl, 25 mM CaCl<sub>2</sub>, 8 mM MgCl<sub>2</sub>, 10 mM Tris and 11 mM Glucose, adjusted to pH 7.4 with NaOH. All chemicals used in the preparation were purchased from Sigma-Aldrich.

**Recording of the electromechanical action potential.** The electrical signal was recorded using a Powerlab 8/35 (ADInstruments Europe, Oxford, UK). It was preamplified with a differential amplifier DP-304 from Warner Instrument Corporation with a gain x1000 using a 3 kHz lowpass and a 10 Hz highpass filter. The mechanical displacement was obtained using an AFM, NanoWizard II from JPK (Germany). The signal was fed into the PowerLab 8/35 and analyzed simultaneously with the electrical recording without any preamplification. The AFM was mounted on the top of an inverted microscope Olympus IX71 (Olympus Corporation, Japan) placed on an anti-vibration table TS-150 (low power, from Herzan LLC, USA). The full setup containing the AFM, optical microscope as well as the anti-vibration table was placed inside a Faraday cage in order to avoid external electrical noise. We used tipless cantilevers from Mikromasch Europe (type HQ:CSC37 and HQ:CSC38). The resonance frequency is 20 kHz for the HQ:CSC37 cantilever and the force constant is 0.3 N/m. The resonance frequency is 10 kHz for the HQ:CSC38 cantilever and the force constant is 0.03 N/m. Thus, the resonance frequencies are outside of the range of the expected signal (1–2  $\mu$ s). Both cantilevers were used under the same experimental conditions. The recording signal frequency was 40 kHz for both, the electrical and mechanical measurement. All the experiments were performed at room temperature of about 22 C.

**Nerve chamber.** We used two different nerve chambers. For the collision experiment the nerve chamber is composed by an array of 21 stainless steel electrodes in a longitudinal cavity covered by a lid in order to isolate the nerve once extracted. The lid also allows to keep the nerve in a saturated vapor atmosphere to prevent the ventral cord from drying. The nerve chamber is a 7 x 2.5 cm block by 1 cm height made on Plexiglass that contains two longitudinal perforation of 1.5 cm length by 0.5 cm wide and 0.5 cm width (19). The large longitudinal aperture contains an array of 21 perforations to allocate the stainless steel electrodes. The array was placed about 0.25 cm from the top of the chamber. The distance between two consecutive electrodes is of 0.25 cm. The stainless steel electrodes have a length of about 3.4 cm and a diameter of 0.5 mm and were fixed in the perforation along the chamber by using a rubber replica casting system (Reprorubber, Islandia, NY) composed by a base and a catalyst from Flexbar Machine Corp (Islandia, NY). For the AFM experiment we used a variation of the previous chamber with half size thickness and with a small open area in the middle without electrodes that allows access to the AFM cantilever.

**Sample preparation.** The Lobster, *Homarus americanus*, was anesthetized by keeping the animal in the freezer for about 30 min. Once removed from the freezer the animal was placed

on the dissecting table and the legs were removed. A second cut was made at the beginning of the abdomen to separate the abdominal part from the thorax. In the thorax, we removed the carapace and made the extraction dorsally. The ventral cord is contained in a tube-like structure formed by the exoskeleton of the lobster and is easy to remove after breaking the shell with scissors. The thoracic segment of the ventral cord including the brain was extracted as a single piece. Special care was taken to remove the circumesophageal connectives from their location surrounding the esophagus. The brain, circumesophageal connectives and subesophageal ganglia was removed from the rest of the ventral cord as placed in a petri-dish with lobster saline solution. The basic anatomical features as well as the basic steps in the preparation are described in the literature (29).

The two circumesophageal connectives were severed at the level of the brain and the subesophageal ganglia (Fig. 2A). From each circumesophageal connective we remove the external connective sheath to expose the main median giant axon (with diameters between 150 and 200  $\mu\text{m}$ , cf. Fig. 2B and C). The preparation was transferred to the AFM nerve chamber with the medial giant axon facing the AFM cantilever (Fig. 2D). For the collision experiment the external connective layer was not removed ensuring a longer survival of the nerve. One of the giant axons was cut at one or two position to rule out the possibility that propagation of pulses in opposite directions occurs in different axons.

**Collision experiments.** The collision experiment was performed independently using a PowerLab 26T from ADInstruments. The instrument possesses an internal bio-amplifier that allows the recording of small electrical potential on the order of microvolts. The bio-amplifier contains two recording channels (further description see ADInstruments webpage). We used the Labchart software from ADInstruments in order to record the signals coming from the ventral cord. The recording frequency was 40 kHz. All experiments were performed at room temperature of about 22 C.

In some experiments we cut the LG axon of the connective at two locations in order to observe a collision of pulses in the MG axon only (cf. Fig. 4, right). In order to make sure that the same axon was cut twice, we performed the following tests:

1. The intact connective (before removing the sheath and cutting the LG axon), we observed two signals from each side corresponding to the action potentials in the MG and LG axons.
2. We performed on single cut at the one end of LG axon. Then, connective was tested again by stimulating at one end of axon before the cut and recording after cut. Only a single peak originating from the MG axon was observed. When placing the stimulation electrode behind the cut, two peaks from the LG and MG axons could be observed.
3. A second cut was made close to the other end of connective. When placing the stimulation electrodes before the first cut and the recording electrodes after the second cut, only one action potential originating from the MG axon was observed, while LG signal was absent.

**Ethical.** The work described in this article has been carried out in accordance with the policy on the use of animals of the Society for Neuroscience.